Asymmetric sulfoxidation of thioether catalyzed by soybean pod peroxidase to form enantiopure sulfoxide in water-in-oil microemulsions: kinetic model

yuanyuan Zhang¹, Huiling Li¹, Zhiyong Wang¹, Depeng Li¹, and Xin Xin Gao¹

¹Qingdao University of Science and Technology

November 24, 2020

Abstract

The asymmetric sulfoxidation catalyzed by soybean pod peroxidase (SPP) in water-in-oil microemulsions were carried out with the yield of 91.56% and e.e of 96.08% at the activity of SPP of 3200 U ml⁻¹ and 50 for 5 h. The mechanism with a two-electron reduction of SPP-I is accompanied with a single-electron transfer to SPP-I and nonenzymatic reactions, indicating that three concomitant sub-mechanisms contribute to the asymmetric oxidation involving five enzymatic and two nonenzymatic reactions, which can represent the asymmetric sulfoxidation of organic sulfides to form enantiopure sulfoxides. With 5.44% of the average relative deviation, a kinetic model fitting experimental data very well was developed. The enzymatic reactions may follow ping-pong mechanism with substrate inhibition of H₂O₂ and product inhibition of esomeprazole, while nonenzymatic reactions, a power law. Those results indicate that SPP with a lower cost and higher thermal stability may be used as an effective substitute for Horseradish Peroxidase.

Hosted file

manuscripts-BB.pdf available at <https://authorea.com/users/364771/articles/494897-asymmetric-sulfoxidation-of-thioether-catalyzed-by-soybean-pod-peroxidase-to-form-enantiopure-sulfoxide-in-water-in-oil-microemulsions-kinetic-model>